

The Genetics of Cream Coat Color in Dogs

SHEILA M. SCHMUTZ AND TOM G. BERRYERE

From the Department of Animal and Poultry Science, University of Saskatchewan, Saskatoon, Canada S7N 5A8.

Address correspondence to S. M. Schmutz at the address above, or e-mail: sheila.schmutz@usask.ca.

Abstract

Cream dogs of several breeds require a genotype of *e/e* at *MC1R* based on 27 individuals in this study. All Akita, Caucasian Mountain Dogs, German Shepherd Dogs, Miniature Schnauzer, and Puli with this genotype are cream, suggesting they are fixed at a second locus which causes the phaeomelanin pigmentation caused by this genotype to be diluted or pale. Conversely, although all Chinese Shar-Pei and Poodles that were cream had an *e/e* genotype at *MC1R*, not all dogs with this genotype are cream. Today, many Golden Retrievers and Labrador Retrievers with an *e/e* genotype are cream instead of the traditional yellow to golden color seen in the past. The second gene in these breeds must have multiple alleles, only one of which causes phaeomelanin pigment to be diluted or pale. *Tyrosinase* (*TYR*) and *solute carrier family 45, member 2* (*SLC45A2*) have been shown to cause cream coat color in other species and were therefore investigated in dogs as candidate genes for this second locus. Although polymorphisms were detected in cDNA sequence from *TYR* and *SLC45A2*, no polymorphism was consistently associated with cream dogs or cosegregated with cream coat color in any of the families used in this study. A microsatellite was detected in a published BAC sequence (GenBank no. AAEX01017083) in intron 2 and was used to map *SLC45A2* to CFA4.

Cream or white is a relatively common coat color in many breeds but the underlying genetic mechanism for this color has not been elucidated. All Samoyed, American Eskimo Dogs, and West Highland White Terriers are white. In recent years, many Labrador Retrievers, of the type that were known as “yellow labs,” are often more cream than yellow (Figure 2 color plate, bottom right corner). Likewise some Golden Retrievers are more cream than golden.

Cream to white also occurs in many other breeds as one of several coat colors. In some breeds, such as German Shepherd Dogs, this coat color was excluded from the show ring but a small group of aficionados has embraced these dogs and formed a separate club.

Little (1957) suggested that an allele e^{cb} of the C locus would pale phaeomelanin to cream and that another possible allele e^l might dilute phaeomelanin to white in addition to the e^l allele which causes albinism in homozygotes. In several species of animals the C locus is considered to be tyrosinase because albinism is caused by mutations at this locus in mice (Yokoyama et al. 1990), humans (Oetting and King 1994; Fukai et al. 1995), rabbits (Aigner et al. 2000), cattle (Schmutz et al. 2004), and cats (Schmidt-Kuntzel et al. 2005; Imes et al. 2006).

Solute carrier family 45, member 2 (*SLC45A2*), previously called *MATP* or *AIM-1*, has been shown to be the gene causing the chestnut/palmino/buckskin and bay/cremello/perlino coat colors in the horse (Locke et al. 2001, Mariat

et al. 2003), underwhite in the mouse (Sweet et al. 1998), a gold color in Medaka fish (Fukamachi et al. 2001), and oculocutaneous albinism type 4 in humans (Newton et al. 2001). There appears to be a codominant dilution of phaeomelanin pigmentation resulting in red, apricot, or cream in some dogs (Sponenberg and Rothschild 2001), such as poodles, with *MC1R* *e/e* genotypes (Newton et al. 2000) similar to the chestnut, palmino and cremello colors in horses (Locke et al. 2001; Mariat et al. 2003), and the Uw^{dbr} allele in mice (Sweet et al. 1998).

We therefore studied *TYR* and *SLC45A2* in the dog as candidate genes for cream coat color in some breeds of dogs in which dogs of *e/e* genotype at *MC1R* varied in shade of phaeomelanin pigmentation, such as Labrador Retrievers, Golden Retrievers, and Chinese Shar-Pei. We also genotyped several cream to white coat color dogs, and their family members when available, for *MC1R* alleles to determine if only phaeomelanin was expressed (*e/e* genotype) (Newton et al. 2000) or if both eumelanin and phaeomelanin were expressed (*E* or E^M allele present) (Schmutz et al. 2003).

Materials and Methods

Dogs and Families

Cheek brush DNA samples (Epicentre, Madison, WI) from 11 dog families that segregated for cream coat color were

Table 1. Primers and the fragment of the dog *SLC45A2* gene amplified

Fragment detected	Direction	Sequence
Exon 1 SNP	Forward	ATGACCACCTTTGATCC TGTGGAG
Exon 1 SNP	Reverse	CTAGGAGAGACAATC CGTTC
Intron 2 microsatellite	Forward	CCACCCTTGCTTCTGT GCTC
Intron 2 microsatellite	Reverse	GTTAAATGAGGTCAT GAGGG
cDNA (5' UTR–Exon 2)	Forward	GACCATCTCTGTTGG CTGCTCAG
cDNA (5' UTR–Exon 2)	Reverse	GGTAGTGGAGGCCCC TCTCC
cDNA (exons 2–5)	Forward	CATTAAAGCCTACTTA TTTG
cDNA (exons 2–5)	Reverse	GATGCACAAGCCCCA ACAT
cDNA (exon 5–3' UTR)	Forward	GTGCACACAACTCCA CAGAG
cDNA (Exon 5–3' UTR)	Reverse	GTAGGGACAGTGTCT CTTTATTG

obtained from dog breeders and owners and used for genotyping. These included Akita (1), Caucasian Mountain Dog (1), Chinese Shar-Pei (2), Golden Retriever (1), Labrador Retriever (1), Poodle (3), and Puli (2) families. Photographs were also supplied to document coat color. Additional dog families were used for linkage mapping. In addition, 27 cream- to white-colored individual dogs of several breeds were also available through the course of our collection of various coat color studies.


We used cDNA prepared from skin biopsies, tail dockings, and surgeries collected from previous coat color studies (Schmutz et al. 2002, 2003) and placed in liquid nitrogen or RNAlater (Ambion Inc., Austin, TX) within 20 min of collection to obtain RNA sequence of *TYR* and *SLC45A2*.

Polymorphism Detection

MC1R genotypes were obtained from all dogs in the study using the PCR–RFLP tests to detect the E^M , E , and e alleles reported previously (Schmutz et al. 2003).

A *TYR* polymorphism was detected as previously reported (Schmidtz and Schmutz 2002). Microsatellite alleles of *FH2312* were detected after polyacrylamide electrophoresis (Mellersh et al. 2000).

In order to study the cosegregation of these dilute phenotypes in these dog families, we tried to identify polymorphisms to use as markers in cosegregation analysis. *SLC45A2* dog sequence was obtained using primers (Table 1) that were initially designed from a dog BAC sequence (GenBank no. AAEX01017083) identified by using a blast search of human *SLC45A2* sequence (GenBank no. NM_01618). The PCR for exons 1 and 2 of *SLC45A2* consisted of 15 μ l which contained 50–100 ng of DNA template, 1.5 μ l of 10 \times PCR buffer (Invitrogen Co., Carlsbad, CA), 1–3 mM MgCl₂

**Figure 1.** Pedigree of an Akita Family that segregated for cream coat color (open symbols) illustrating that all the cream pups inherited an e/e *MC1R* genotype.

(Invitrogen), 0.2 mM dNTP (Invitrogen), 0.5 U *Taq* DNA polymerase (Invitrogen), 0.66 pmol of each primer, and 9.6 μ l of ddH₂O. The reaction began with an initial 4-min denaturation step at 94 °C; followed by 35 cycles of 50 s at 94 °C, 50 s at 57 °C, and 50 s at 72 °C; and finished with a final 4-min extension step at 72 °C.

Two polymorphisms were identified in *SLC45A2*. A single-nucleotide polymorphism (SNP) (150G>A) (GenBank no. DQ118774) was identified in exon 1 of *SLC45A2* that did not alter the amino acid. A natural *AaI* cut site occurs at this SNP. The A allele cuts into 226- and 248-bp fragments. The G allele cuts into 93, 115, and 226-bp fragments that can be resolved on 2% agarose gel.

A microsatellite in intron 2 of *SLC45A2* was identified in the BAC sequence (GenBank no. AAEX01017083). In order to detect this microsatellite, the forward primer was end-labeled in a 10- μ l reaction consisting of 2.0 μ l of forward primer, 5.0 μ l dH₂O, 1.0 μ l of 10 \times polynucleotide kinase (PNK) buffer, 1 μ l ATP³², and 1 μ l of PNK (New England BioLabs Inc., Beverly, MA). This reaction was then incubated at 37 °C for 30 min. The PCR protocol was the same as for the exon 1 SNP with the exception of the annealing temperature being 55 °C. The alleles were differentiated on a 6% polyacrylamide gel and visualized on autoradiograph film. Seven alleles were identified among the 35 dogs genotyped for the microsatellite in intron 2. The alleles ranged in size from 220 to 296 bp.

Results and Discussion

In the Akita (Figure 1), Caucasian Mountain Dog, both Chinese Shar-Pei, all three Poodle, and both Puli families, the cream-colored pups had an e/e genotype at *MC1R*, as shown in Figure 2 (Supplemental Table 1). In addition, we also found that 17 individual dogs that were cream to white from breeds, in which cream is one of several coat colors, all had an e/e genotype at *MC1R*. These included 3 Cardigan Welsh Corgi, 3 Chinese Shar-Pei, 5 German Shepherd Dogs, 2 Great Pyrenees, and 3 Miniature Schnauzer. Furthermore white dogs of other breeds where this is the only color, such as American Eskimo Dog (6), Samoyed (3), and West Highland White Terrier (1), also genotyped as e/e at *MC1R*. In



Figure 2. Photographs of representative cream dogs that had a genotype of e/e at *MC1R*. Top to bottom: Akita pup, Miniature Schnauzer, German Shepherd Dog, Puli, and Caucasian Mountain Dog. The Labrador Retriever littermates in the final photo illustrate the difference between cream or pale yellow and dark yellow seen in some breeds.

both the Golden Retriever and yellow Labrador Retriever families, although all individuals had an e/e genotype at *MC1R*, some pups were golden/yellow and others were cream (Figure 2, bottom right).

In all the families studied, cream coat color fit an autosomal recessive inheritance pattern among the dogs with e/e *MC1R* genotype. This e/e genotype alone does not cause cream in all dogs of all breeds, and so we assumed that some other gene, which varied in some breeds, must interact to cause cream instead of yellow or red.

TYR

We obtained the complete coding sequence of *TYR* (GenBank no. AY336053) from a cream Poodle, an albino Lhasa Apso, and 2 albino Doberman Pinschers as well as a blue Doberman Pinscher and black-and-white Large Munsterlanders as controls. Although several SNPs were detected (780A>T, 882A>G, 1223G>A which caused an arginine to be replaced by a glutamine, 1312C>T which caused a lysine to be replaced by a phenylalanine), none were consistent




Figure 3. Pedigree of a Golden Retriever Family that segregated for cream coat color (open symbols) illustrating the paternally and maternally inherited alleles of *FH2312* alleles, a microsatellite near *TYR*. All Golden Retrievers have an *e/e* *MC1R* genotype.

with either cream or albinism when compared with the sequence obtained from additional dogs of the same phenotype.

We had previously identified a polymorphism in *TYR*, 175G>A (GenBank no. AF473807), which changed a valine to an isoleucine (Schmidt and Schmutz 2002) that was used to map this gene to CFA21 with no recombinants to *FH2312*. Some cream dogs were heterozygous, suggesting that this variant was not responsible for cream coat color which is inherited as a recessive. Neither the informative Golden Retriever family of 7 pups (Figure 3) nor the Labrador Retriever family of 4 pups showed cosegregation with cream coat color and *TYR* using the *FH2312* microsatellite (LOD = -5.02). This also excludes mutations in the promoter and introns as being associated with variation in phaeomelanin pigmentation.

SLC45A2

No mutations affecting amino acids were found in the coding sequence of *SLC45A2* (GenBank no. DQ302162) in a cream poodle compared with a Brittany Spaniel of darker red pigmentation and *e/e* genotype at *MC1R* or a black-and-white Large Munsterlander of *E/E* genotype. We did identify an SNP, 150G>A, in exon 1 (GenBank no. DQ118774S1), but it was not informative in the families segregating for shades of phaeomelanin. We also identified a microsatellite in intron 2. This did not cosegregate with pale and dark yellow in the Labrador Retriever family (Figure 4) (LOD = -2.8).

This microsatellite was polymorphic in several additional dog families. Linkage mapping was accomplished using one French Brittany, one Newfoundland, one Cardigan Welsh Corgi, and one Tervuren family. *SLC45A2* was mapped to canine chromosome 4.9 cm from *FH2097* (LOD = 5.158).

Because cream dogs always have an *e/e* genotype at *MC1R*, DNA testing for an *e* allele should be predictive that the dog is heterozygous for cream coat color in breeds such as Akita, Caucasian Mountain Dogs, German Shepherd Dogs, Miniature Schnauzers, and Puli. Neither *TYR* nor *SLC45A2* appeared to co-segregate with cream coat color in dogs from breeds with an *e/e* genotype where the variation




Figure 4. Pedigree of a Labrador Retriever Family that segregated for pale (open symbols) and dark yellow among the yellow pups which all have an *e/e* *MC1R* genotype. The paternally and maternally inherited alleles of the microsatellite detected in intron 2 of *SLC45A2*.

in phaeomelanin can be cream through yellow and even red, such as Chinese Shar-Pei, Golden Retrievers, Labrador Retrievers, and Poodles.

Supplementary Material

Supplementary Table 1 can be found at <http://www.jhered.oxfordjournals.org/>.

Acknowledgments

We thank HealthGene, Inc., and a Collaborative Research and Development matching grant from the Natural Science and Engineering Research Council for funding. We are grateful to the many dog owners who contributed cheek brush and/or skin samples from their dogs. Our thanks to the very helpful suggestions of 2 anonymous reviewers.

References

- Aigner B, Besenfelder U, Müller M, Brem G. 2000. Tyrosinase gene variants in different rabbit strains. *Mamm Genome*. 11:700–702.
- Fukai K, Holmes SA, Lucchese NJ, Mok Siu V, Weleber RG, Schnur RE, Spritz RA. 1995. Autosomal recessive ocular albinism associated with a functionally significant tyrosinase gene polymorphism. *Nat Genet*. 9:92–95.
- Fukamachi S, Shimada A, Shima A. 2001. Mutations in the gene encoding B, a novel transporter protein, reduce melanin content in medaka. *Nat Genet*. 28:381–385.
- Imes DL, Geary LA, Grahm RA, Lyons LA. 2006. Albinism in the domestic cat (*Felis catus*) is associated with a tyrosinase (*TYR*) mutation. *Anim Genet*. 37:175–178.
- Little CC. 1957. *The inheritance of coat color in dogs*. Ithaca (NY): Comstock.
- Locke MM, Ruth LS, Millon LV, Penedo MC, Murray JD, Bowling AT. 2001. The cream dilution gene, responsible for the palomino and buckskin coat colours, maps to chromosome 21. *Anim Genet*. 32:340–343.
- Mariat D, Taourit S, Guerin G. 2003. A mutation in the *MATP* gene causes the cream coat colour in the horse. *Genet Sel Evol*. 35:1–16.
- Mellersh C, Hitte C, Richman M, Vignaux F, Priat C, Jouquand S, Werner P, André C, DeRose S, Patterson D, et al. 2000. An integrated linkage-radiation map of the canine genome. *Mamm Genome*. 11:120–130.

- Newton J, Wilkie A, He L, Jordan S, Metallinos D, Holmes N, Jackson IJ, Barsh GS. 2000. Melanocortin 1 receptor variation in the domestic dog. *Mamm Genome*. 11:24–30.
- Newton JM, Cohen-Barak O, Hagiwara N, Gardner JM, Davisson MT, King RA, Brilliant MH. 2001. Mutations in the human form of oculocutaneous albinism, OCA4. *Am J Hum Genet*. 69:981–988.
- Oetting WS, King RA. 1994. Analysis of tyrosinase mutations associated with tyrosinase-related oculocutaneous albinism (OCA1). *Pigment Cell Res*. 7:285–290.
- Schmidt-Kuntzel A, Eizirik E, O'Brien SJ, Menotti-Raymond M. 2005. Tyrosinase and tyrosinase related protein 1 alleles specify domestic cat coat color phenotypes of the albino and brown loci. *J Hered*. 96:289–96301.
- Schmidtz BH, Schmutz SM. 2002. Linkage mapping of *TYR* to dog chromosome 21. *Anim Genet*. 33:476–477.
- Schmutz SM, Berryere TG, Ciobanu DC, Mileham AJ, Schmidtz BH, Fredholm M. 2004. A form of albinism in cattle is caused by a tyrosinase frame-shift mutation. *Mamm Genome*. 15:62–67.
- Schmutz SM, Berryere TG, Ellinwood NM, Kerns JS, Barsh GS. 2003. *MC1R* studies in dogs with melanistic mask or brindle patterns. *J Hered*. 94:69–73.
- Schmutz SM, Berryere TG, Goldfinch AD. 2002. *TYRP1* and *MC1R* genotypes and their effects on coat color in dogs. *Mamm Genome*. 13:380–387.
- Sponenberg DP, Rothschild MF. 2001. Genetics of coat colour and hair texture In: Ruvinsky A, editor. *Genetics of the dog*. New York: CABI Press.
- Sweet HO, Brilliant MH, Cook KR, Davisson MT. 1998. A new allelic series for the underwhite gene on mouse chromosome 15. *J Hered*. 89:546–551.
- Yokoyama T, Silversides DW, Waymire KG, Kwon BS, Takeuchi T, Overbeek PA. 1990. Conserved cysteine to serine mutation in tyrosinase is responsible for the classical albino mutation in laboratory mice. *Nucleic Acids Res*. 18:7293–7298.

Corresponding Editor: Urs Giger